



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12Q 1/68	A1	(11) International Publication Number: WO 96/31622 (43) International Publication Date: 10 October 1996 (10.10.96)
(21) International Application Number: PCT/GB96/00848 (22) International Filing Date: 9 April 1996 (09.04.96) (30) Priority Data: 9507238.5 7 April 1995 (07.04.95) GB (71) Applicant (for all designated States except US): ISIS INNOVATION LIMITED [GB/GB]; 2 South Parks Road, Oxford OX1 3UB (GB). (72) Inventors; and (75) Inventors/Applicants (for US only): SOUTHERN, Edwin, Mellor [GB/GB]; 12 School Road, Kidlington OX5 2HB (GB). PRITCHARD, Clare, Elizabeth [GB/GB]; 43 Cherry Tree Close, Southmoor, Abingdon OX13 5BE (GB). CASE-GREEN, Stephen, Charles [GB/GB]; Flat 2, Sylvia House, 32A Union Street, Oxford OX4 1JP (GB). (74) Agent: PENNANT, Pyers, Stevens Hewlett & Perkins, 1 Serjeants' Inn, Fleet Street, London EC4Y 1LL (GB).		(81) Designated States: JP, US, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: DETECTING DNA SEQUENCE VARIATIONS (57) Abstract <p>A method of analysing a polynucleotide target involves incubating the target with an oligonucleotide probe, generally an array of immobilised oligonucleotide probes, to form a duplex, and using ligase or polymerase to extend one chain of the duplex. A point mutation or variable number tandem repeat section may be analysed. Arrays of immobilised oligonucleotides are provided for use in the method.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

DETECTING DNA SEQUENCE VARIATIONS

5 INTRODUCTION

Detection of variation in DNA sequences forms the basis of many applications in modern genetic analysis: it is used in linkage analysis to track disease genes in human pedigrees or economically important traits in animal and plant breeding programmes; it forms the basis of

10 fingerprinting methods used in forensic and paternity testing [Krawczak and Schmidtke, 1994]; it is used to discover mutations in biologically and clinically important genes [Cooper and Krawczak, 1989]. The importance of DNA polymorphism is underlined by the large number of methods that have been developed to detect and measure it [Cotton, 1993]. Most of

15 these methods depend on one of two analytical procedures, gel electrophoresis or molecular reassociation, to detect sequence variation. Each of these powerful procedures has its drawbacks. Gel electrophoresis has very high resolving power, and is especially useful for the detection of variation in the mini- and microsatellite markers that are used in linkage

20 analysis and fingerprinting; it is also the method used to analyse the variation found in the triplet repeats that cause a number of mutations now known to be the cause of around ten genetic disorders in humans [Willems, 1994]. Despite its great success and widespread use, gel electrophoresis has proved difficult to automate: even the systems which

25 automate data collection require manual gel preparation; and as samples are loaded by hand, it is easy to confuse samples. The continuous reading electrophoresis machines are expensive, and manual analysis is technically demanding, so that its use is confined to specialised laboratories which have a high throughput. Furthermore, difficulties in

30 measuring fragment size preclude rigorous statistical analysis of the

results.

By contrast, oligonucleotide hybridisation lends itself to automation and to quantitative analysis [Southern *et al.*, 1992], but it is not well suited to the analysis of variation in the number of repeats in the micro- and minisatellites, as the small fractional change in the number of repeats produces a barely detectable change in signal strength; and of course it would not be possible to distinguish two alleles in the same sample as each would contribute to a single intensity measurement. Thus, many different combinations of alleles would produce the same signal.

Present hybridisation methods are much better suited to analysing variation in the DNA due to point mutation - base substitution deletions and insertions, for which it is possible to design allele specific oligonucleotides (ASOs) that recognise both the wild type and the mutant sequences [Conner *et al.*, 1983]. Thus it is possible in principle, in a relatively simple test, to detect all possible genotypes. However, a problem that arises in practice in the use of oligonucleotide hybridisation is that in some cases the extent of reassociation is barely affected by a mismatched base pair.

THE INVENTION

The invention describes a general approach which can be applied to all forms of variation commonly used as DNA markers for genetic analysis. It combines sequence-specific hybridisation to oligonucleotides, which in the preferred embodiment are tethered to a solid support, with enzymatic reactions which enhance the discrimination between matching and non-matching duplexes, and at the same time provide a way of attaching a label to indicate when or which reaction has taken place. Two enzymatic reactions, chain extension by DNA dependent DNA polymerases and DNA strand-joining by DNA ligases, are dependent on perfect matching of sequences at or around the point of extension or joining. As we shall show, there are several ways in which these enzymes

can be used with sequence-specific oligonucleotides to detect variation in target sequences.

In all cases, the sequence to be analysed, the target sequence, will be available as a nucleic acid molecule, and may be a DNA molecule produced, for example, by the polymerase chain reaction. However, the methods are not confined to analysis of DNA produced in this way. In all applications, the target sequence is first captured by hybridisation to oligonucleotides which are preferably tethered to a solid support; for example, the oligonucleotides may be synthesised *in situ* as described [Maskos and Southern, 1992]; or they may be presynthesised and then coupled to the surface [Khrapko *et al.*, 1991].

In one aspect of the invention the novelty arises from the exploitation of enzymes in combination with substrates or primers tethered to solid supports. A further novelty exploits the observation that DNA ligases and polymerases can be used to distinguish sequence variants which differ in the number of units of a tandemly repeating sequence. This observation is surprising, as tandemly repeated sequences can form duplex in any register, thus in principle, length variants can form duplexes which match at the ends even when the two strands contain different numbers of repeat units. Although we demonstrate the application of this method in conjunction with tethered oligonucleotides, it should be evident that this reaction could be used to analyse VNTR (variable number tandem repeat) sequences in the liquid phase followed by some other method of analysis, such as gel electrophoresis.

In one aspect the invention provides a method of analysis which comprises: providing a polynucleotide target including a nucleotide at a specified position, and an oligonucleotide probe, tethered to a support, said probe being complementary to the target and terminating at or close to the said specified position; and performing the steps:

- a) incubating the target with the probe to form a duplex,

- b) incubating the duplex under ligation conditions with a labelled oligonucleotide complementary to the target,
- c) and monitoring ligation in b) as an indication of a point mutation at the specified position in the target.

5 In another aspect the invention provides a method of analysis which comprises: providing a polynucleotide target having a variable number tandem repeat section and a flanking section, and an oligonucleotide probe having a section complementary to the repeat section and a flanking section of the target; and performing the steps:

- 10 a) incubating the target with the probe to form a duplex,
- b) incubating the duplex with a labelled oligonucleotide and/or at least one labelled nucleotide under chain extension conditions,
- c) and monitoring chain extension as an indication of the length of the variable number repeat section of the target.

15 A polynucleotide target is provided, in solution when the probe is tethered to a support, and may be DNA or RNA. This polynucleotide target is caused to hybridise with an oligonucleotide probe. The term oligonucleotide is used here, as common terminology for the primers and substrates commonly utilised by polymerase and ligase
20 enzymes. However, the term is used in a broad sense to cover any substance that serves as a substrate for the enzymes, including single stranded chains of short or moderate length composed of the residues of nucleotides or of nucleotide analogues, and also longer chains that would ordinarily be referred to as polynucleotides.

25 The probe may be tethered to a support, preferably by a covalent linkage and preferably through a 5' or 3' terminal nucleotide residue. An array of oligonucleotide probes may be tethered at spaced locations, for example on a derivatised glass surface or the surface of a silicon microchip, or alternatively on individual beads.

30 In another aspect the invention provides an array of

oligonucleotides, for analysing a polynucleotide target containing a variable sequence, in which each component oligonucleotide i) comprises a sequence complementary to the target including an expected variant of the target, and ii) is tethered to a solid support in a chemical orientation which
5 a) permits duplex formation with the target, and b) permits chain extension only when the sequence of the oligonucleotide matches the variable sequence of the target.

In another aspect the invention provides a set or array of oligonucleotides, for analysing a polynucleotide target containing a variable
10 number tandem repeat sequence, in which each component oligonucleotide i) comprises a sequence complementary to a part of the target immediately adjacent the repeat sequence, ii) comprises a sequence complementary to the repeat sequence of the target and containing a number of repeats expected in the target, and iii) is configured
15 in a way that a) permits duplex formation with the target, and b) permits chain extension only when the number of repeats in the oligonucleotide equals or is less than the number of repeats in the target.

In another aspect the invention provides an array of oligonucleotides in which different oligonucleotides occupy different
20 locations and each oligonucleotide has a 3' nucleotide residue through which it is covalently tethered to a support and a 5' nucleotide residue which is phosphorylated.

The invention also provides a method of making an array of different oligonucleotides tethered to different locations of a support, which
25 method comprises the steps of: providing a first intermediate oligonucleotide tethered to the support and a second intermediate oligonucleotide in solution, and a third oligonucleotide that is complementary to both the first and second intermediate oligonucleotides, forming a duplex of the third oligonucleotide with the first and second
30 intermediate oligonucleotides, and ligating the first intermediate

oligonucleotide with the second intermediate oligonucleotide; and repeating the steps with oligonucleotides tethered to different locations of the support

Reference is directed to the accompanying drawings in which
5 each of Figures 1 to 6 is a series of diagrams illustrating a method according to the invention.

Figure 1 shows detection of point mutation by single base extension.

Figure 2 shows detection of point mutation by hybridisation to
10 allele specific oligonucleotides and chain extension.

Figure 3A shows detection of point mutation by tag ligation to allele specific oligonucleotides.

Figure 3B shows detection of point mutation by ligation to library of differentially tagged allele specific oligonucleotides.

15 Figure 4A shows analysis of variable numbered tandem repeats by ligation of tag to allelic variants.

Figure 4B shows analysis of variable number tandem repeats by ligation of tag to allelic variants.

Figure 5 shows measurement of variable number tandem
20 repeats by labelled chain extension.

Figure 6 shows analysis of variable number tandem repeats by ligation of tag followed by chain extension.

DETAILED DESCRIPTION

25 Detection of point mutation

I. Single base-specific extension of tethered primers.

In this application, the tethered oligonucleotide terminates at a position one base before the variable base in the target sequence (Fig.1). A nucleotide precursor triphosphate or dideoxynucleotide
30 triphosphate, labelled, for example with a fluorescent tag, is added in the

presence of a nucleic acid synthesising enzyme which requires a specific template in order to incorporate the complementary base. In the case of DNA polymerase, the labelled base will be incorporated from a deoxyribonucleotide precursor only if the precursor base is complementary to the base in the target sequence. Thus, mutants will give a negative result.

II. Chain extension from tethered ASOs.

In this case, the tethered oligonucleotide terminates in a base which is complementary to the variable base in the target sequence. Labelled precursor nucleoside triphosphates and polymerase are added. Polymerisation takes place only if the last base of the primer is complementary to the variable base in the target (Fig.2). Thus, mutants will give a negative result.

III. Ligation of tag sequences to tethered ASOs.

In this method, the tethered oligonucleotide may end at the variable position in the target sequence, or it may end close to this position. In either case, hybridisation of the target to the tethered ASOs will produce a substrate for ligating a tag oligonucleotide only if the bases at the join are well matched (Fig. 3a). Thus, mutants which are close enough to the joining position to prevent ligation will give a negative result. Alternatively, the tethered oligonucleotide may terminate at the base before the variable position; in this case, the ligation reaction can be carried out using a mixture of tag oligonucleotides, one for each of the possible alternative variants. Each tag would be labelled differently, for example, with a different fluophore, so that those that ligated could be recognised identifying the variant base (Fig. 3b).

Analysis of VNTR Lengths by Ligation to Anchored VNTRS

In this application "tag" oligonucleotides are ligated to sets of tethered oligonucleotides after hybridisation of the target, which acts as a template to bring the tags and the tethered oligonucleotides together.

5 In Figure 4a, the tethered oligonucleotides comprise three parts, an "anchor" sequence which is common to all members of a VNTR set, which is attached to the solid support and which hybridises to a sequence flanking the variable region, a variable number of the repeated sequence unit, and a distal sequence. Each allele, represented by a
10 different number of repeats, is located on a different solid support or at a different location on the same solid surface. Hybridisation of the target sequence will produce a series of duplexes, the structures of which depend on the number of units that the target contains. If the number matches the number in a tethered oligonucleotide, the target will meet the end of a tag
15 when the tag is hybridised to the distal sequence of the tethered oligonucleotide. If the number is greater or smaller, there will be a gap in the duplex which reduces or prevents ligation of the tag.

In Figure 4b, the tethered oligonucleotides comprise two parts: an "anchor" sequence which is common to all members of a VNTR
20 set and which hybridises to a sequence flanking the repeated region, and a variable number of the repeated sequence unit. Each allele, represented by a different number of repeats, is located on a different solid support, or at a different location on the same solid surface. Hybridisation of the target sequence will produce a series of duplexes, the structures of which depend
25 on the number of repeat units that the target contains (Fig.4b). If the number matches the number in a tethered oligonucleotide, the latter will meet the end of the tag when the tag is hybridised to its complement in the target sequence, and form a substrate for ligation of the tag. If the number is greater or smaller, there will be a gap in the duplex which reduces or
30 prevents ligation of the tag.

Analysis of VNTR Lengths by Chain Extension

The number of repeat units in a VNTR may vary within small limits, and in these circumstances, the method of analysis described above, using a ligase, will be appropriate; in other cases, for example the

5 trinucleotide repeats associated with a number of human inherited disorders, the variation may be too large to analyse in this way. For a number of triplet repeats, the variation can be from around 10-50 in the normal chromosome to more than a thousand in the affected individual (Table 1). It is probably unrealistic to measure such large numbers of

10 repeats using the ligase reaction. In these cases, where the difference between the normal and the mutant allele is large, an alternative is to measure approximately the number of repeat units using labelled precursors with a polymerising enzyme. The enzyme may be either a polymerase, such as DNA dependent DNA polymerase, or a ligase. In the

15 former case, the oligonucleotides have to be tethered at their 5' ends to satisfy the requirement for enzymatic extension by the polymerase. The solid support carries an oligonucleotide anchor complementary to a sequence flanking the repeat unit; for example, the sequence can be that of one of the primers used to amplify the test sequence by the PCR. After

20 hybridisation of the test sequence to the anchor, the repeat insert may be copied by a polymerase or a ligase (Fig. 5) incorporating a labelled precursor. The amount of label incorporated is proportional to the number of repeat units. Incomplete hybridisation of the target to the anchor sequence would give a deceptively low measure of the repeat number.

25 This problem can be overcome by standardising the measurement in one of several possible ways. For example, if the target sequence itself is labelled as shown in Fig. 5, the final measurement will be a ratio of two labels: the target and the incorporated precursor.

Alternatively, in the case of a triplet repeat, incorporation will

30 end at the point in the sequence where the missing precursor base is

needed for further extension; where a ligase had been used to polymerise monomers of the basic repeat unit, this will also end at the end of the VNTR insert. At this point a labelled "capping" sequence can be added by ligation. In such cases, the measurement will be the ratio of cap to
5 polymer labels.

Analysis of VNTR Lengths by Combining Ligation and Chain Extension

A more powerful way of analysing VNTRs which may vary in
10 length over a wide range, would be to test first for ligation to a labelled tag oligonucleotide; this would give the results already described for targets with different lengths of repeats: a negative result where the VNTR lengths were longer or shorter in the target than in the tethered oligonucleotides, and positive results where they were the same. Following ligation, which
15 we have shown can be made to go to completion, the different length classes will behave differently as substrates for DNA polymerase. Those targets in which the repeat number is less than that of the tethered probe will not act as substrates (Fig. 6c). Targets which have the same number of repeats as the probes will not be elongated by polymerase, because the
20 ligated tag will block extension (Fig. 6a). The only cases where extension will occur are those for which the targets are longer than the probes (Fig. 6b). If the analysis is done on an array of probes with different numbers of inserts up to a certain limit, there will be a clear indication of the number of repeats in the targets from the ligation results provided they are within the
25 range of sizes represented in the array of probes. If, within the targets, there is one which is longer than this range, it will show up in the polymerase analysis. This test will be especially useful for the triplet repeats associated with the so-called "dynamic mutations", for example, that which is found in the fragile X mutation, where the size range varies
30 from ca. 10-1000. It would be difficult to accommodate all of these size

classes on a single array.

EXPERIMENTAL SUPPORT FOR THE CLAIMS

Properties of DNA Polymerases and Ligases

5 Most DNA polymerases, reverse transcriptases, some DNA
dependent RNA polymerases and ligases can use as substrates one or
more oligonucleotides which are bound to a long DNA strand through
Watson-Crick base pairing. In the case of polymerases, an oligonucleotide
is used as a primer to which the first base in the growing chain is added.
10 In the case of ligases, two oligonucleotides are joined provided that both
are paired to the DNA strand and perfectly matched in the base pairs at or
close to the junction point. It is these properties that make the enzymes
useful for the detection of DNA sequence variation; in particular, the
requirement for specific base pairing at the site of extension or joining
15 complements the sequence discrimination that is already provided by the
Watson-Crick pairing between the oligonucleotide and the target sequence
that is needed to form a stable duplex. Thus, it has been found that
discrimination by hybridisation alone is most sensitive if the variant base(s)
is (are) close to the middle of the oligonucleotide. By contrast, for the
20 enzymes, discrimination is highest if the variant mismatching bases are
close to the end where the extension or join takes place. Together,
hybridisation under stringent conditions and enzymatic extension or joining
provide greater discrimination than either alone, and several methods have
been developed to exploit this combination in systems for genetic analysis
25 [references in cotton, 1993]. The hybridisation and the enzyme reaction
are normally carried out in solution, following which the product is captured
on a solid support, or separated by gel electrophoresis for detection and/or
measurement.

 In one embodiment, the invention described here employs
30 oligonucleotides coupled to a solid surface, so that the advantages of

working in mixed phase are brought to all steps: hybridisation, enzymatic extension or joining, and detection. This provides great sensitivity and convenience. As many different oligonucleotides can be bound to one surface in an array, it enables many different sequences to be analysed together, in a single reaction; this also ensures that all reactions are carried out under identical conditions, making comparisons more reliable.

Support-bound Oligonucleotides

Two different methods have been developed for making oligonucleotides bound to a solid support: they can be synthesised *in situ*, or presynthesised and attached to the support. In either case, it is possible to use the support-bound oligonucleotides in a hybridisation reaction with oligonucleotides in the liquid phase to form duplexes; the excess of oligonucleotide in solution can then be washed away. Hybridisation can be carried out under stringent conditions, so that only well-matched duplexes are stable. When enzymes are to be used, the chemical orientation of the oligonucleotide is important; polymerases add bases to the 3' end of the chain; ligases join oligonucleotides which are phosphorylated at the 5' end to those with a 3'-OH group. Oligonucleotides tethered to the solid substrate through either end can be made *in situ* by using the appropriate phosphoramidite precursors [references in Beaucage and Iyer, 1992]; or presynthesised oligonucleotides can be fixed through appropriate groups at either end. We will demonstrate that oligonucleotides can be phosphorylated at the 5' end *in situ* using ATP and polynucleotide kinase, or they may be phosphorylated chemically [Horn and Urdea, 1986].

Tethered Oligonucleotides as Substrates for DNA Modifying Enzymes

The applications envisaged here require that the oligonucleotides tethered to the solid substrate can take part in reactions catalysed by DNA polymerases and ligases.

DNA polymerase

The M13 sequencing primer -

5'-GTAAACGACGGCCAGT-3' - attached to aminated polypropylene through its 5' end was synthesised as described. A solution of M13 DNA (single-strand, replicative form, 0.1 µl, 200 ng/µl) was applied in two small spots to the surface of the derivatised polypropylene. A solution containing three non-radioactive deoxyribonucleotide triphosphates, dATP, dGTP, TTP (10 µmol each), $\alpha^{32}\text{P}$ -dCTP (10 µCi), Taq DNA polymerase and appropriate salts, was applied over a large area of the polypropylene, including the area where the M13 DNA had been spotted. The polypropylene was incubated at 37°C for 1 hr in a vapour saturated chamber. It was then washed in 1% SDS at 100°C for one minute, and exposed to a storage phosphor screen for one minute and scanned in a phosphorimager. The regions where the DNA had been applied showed a high level of radioactivity, against a low background where no DNA had been applied. This experiment shows that oligonucleotides tethered to a solid support can act as primers for DNA-dependent synthesis by DNA polymerase, as required for applications using this enzyme for mutation detection.

Experiments described below show that both polynucleotide kinase and DNA ligase can be used to modify oligonucleotides tethered to a solid support. There are several ways in which phosphorylated oligonucleotides and the ligase reaction can be used to detect sequence variation.

Methods for Making Arrays of Sequence Variants.**1. Allele specific oligonucleotides for point mutations.**

For the preferred embodiment, it will be necessary to use oligonucleotides tethered to a solid support. The support may take the form of particles, for example, glass spheres, or magnetic beads. In this

case the reactions could be carried out in tubes, or in the wells of a microtitre plate. Methods for both synthesising oligonucleotides and for attaching presynthesised oligonucleotides to these materials are known [Stahl *et al.*, 1988]. Methods for making arrays of ASO's representing point mutations were described in patent application PCT/GB89/00460 and in Maskos and Southern (1993). We also demonstrated how oligonucleotides tethered to a solid support in an array could distinguish mutant from wild type alleles by molecular hybridisation.

For the present invention, the same methods could be used to create oligonucleotide arrays of ASOs, but in order that they can be used as substrates for the enzymes, they need to be modified; for ligation, it may be necessary to phosphorylate the 5' end; for extension by polymerase it will be necessary to attach to oligonucleotides to the solid substrate by their 5' ends.

15

2. Arrays for scanning regions for mutations.

It is often desirable to scan a relatively short region of a gene or genome for point mutations: for example, many different sites are mutated in the CFTR gene to give rise to cystic fibrosis; similarly, the p53 tumour suppressor gene can be mutated at many sites. The large numbers of oligonucleotides needed to examine all potential sites in the sequence can be made by efficient combinatorial methods [Southern *et al.*, 1994]. A modification of the protocol could allow such arrays to be used in conjunction with enzymes to look for mutations at all sites in the target sequence.

25

3. VNTRs.

The most commonly used VNTRs are repeats of very short units, typically mono to tetranucleotides. However, there is another class, the minisatellites, in which the repeat unit is somewhat longer, up to 20 or

30

more nucleotides. The short repeats may be made using chemical synthesis; in the case of inserts with large numbers of repeat units, it would be more economical to use a synthetic route which used the repeat unit as a reactant, rather than building them up one base at a time; such methods have been used to make polynucleotides by chemical synthesis. An attractive alternative would be to build the repeat units by ligating the monomer units; they could be added stepwise, one unit at a time provided a method could be found to block one end to prevent polymerisation; for example the oligonucleotide building block may be terminated by a hydroxyl group, which is then phosphorylated after ligation so that the unit becomes an acceptor for the next one; the monomer may have a phosphate group protected by a cleavable group, such as a photocleavable group, which can be removed after ligation to allow a subsequent ligation [Pillai, 1980]. A second alternative, which would be especially favourable for longer units such the minisatellites, would be to attach either cloned or enzymatically amplified molecules to the solid support. For example, each variant sequence could be amplified by the PCR, using a biotinylated oligonucleotide for one of the primers. The strand starting with this group could then be attached to a streptavidin coated surface, and the other strand removed by melting [Stahl *et al.*, 1988].

EXAMPLE 1

25 DEMONSTRATION OF THE ANALYSIS OF LENGTH POLYMORPHISM BY LIGATION TO AN ARRAY OF VNTRs

An array of VNTRs was made as described in Fig. 4b, in which the anchor sequence was 5'-tgtagtggtgtgatcaaggc-3'. The repeat unit was 5'-cttt-3'; stripes, ca 3 mm wide, of sequence variants of the form: Anchor-Repeat_N, with N = 4-10, were made as stripes on the surface of a

sheet of polypropylene. The synthesis was carried out using 3'-deoxyribophosphoramidites, this chemical orientation produces oligonucleotides tethered through their 3' ends to the polypropylene, and a free 5' hydroxyl group. To create a substrate for ligation, this OH group
5 was phosphorylated by immersing a strip of the polypropylene (3mm x 18mm), carrying the array of oligonucleotides, in 0.5 ml of a solution containing 4mM ATP and 77.6 units of polynucleotide kinase with buffer and Mg⁺⁺ according to the supplier's instructions. The reaction was left for 6 hours at 37°C; the strip was removed and immersed in boiling water to
10 kill the polynucleotide kinase. The target sequences, which are complementary to elements of the array oligonucleotides and to the ligation tag, 5'-Anchor-Repeat₁₀-Tag and 5'-Anchor-Repeat₅-Tag were added to 0.5 ml of a solution, preheated to 95°C, containing the tag, 5'-gtggctactaaagttctgct-3', which had been labelled at its 5' end using
15 polynucleotide kinase and ³³P-gamma-ATP, thermal ligase (500 units), and buffer and salts according to the supplier's instructions. The polypropylene strip was immersed in the hot solution, which was then allowed to cool to 68°C, and left at this temperature for 16 hours. The polypropylene strip was removed and placed in 25% formamide at 95°C for 5 minutes, rinsed
20 in water at the same temperature, dried and exposed to a storage phosphor screen, from which an image of the radioactivity was collected. The results showed counts close to background over most of the array; counts on Anchor-Repeat₅ and Anchor-Repeat₁₀ were more than five times those over adjacent cells in the array. This experiment indicates that the
25 ligase is able to distinguish length variants of the repeat sequence and gives optimum ligation only when the number of repeats in the target matches that in the allele specific oligonucleotide in the array. Thus, it should easily be possible to detect the two allelic variants in a heterozygote.

30 Ligation and/or polymerisation is possible if the

oligonucleotide is tethered through the 5' end . The oligonucleotides can be synthesised *in situ* using deoxyribophosphoramidites with 3' dimethoxytrityl groups and 5' phosphoramidite (reverse phosphoramidites). It is unnecessary to phosphorylate the tethered polynucleotide for ligation assays since the phosphate needed for ligation is provided by the tag oligonucleotide.

EXAMPLE 2

Analysis of VNTR lengths

10 An array of VNTR's was made as described in Figure 4B, with the oligonucleotides anchored through their 5' ends. The repeat unit was 5' ttca and the anchoring sequence 5' cttatttcctca. Stripes 6mm wide of sequence variants of the form: Anchor-Repeat_N where N= 4-8 were made on the surface of a sheet of polypropylene using "reverse" 15 phosphoramidite monomers.

Analysis by ligation

A strip of the array (30mm x 2mm) was immersed in a solution of 600pmols of the target oligonucleotide 5' 20 cacagactccatgg(tgaa)₆tgagggaaataag, 1.4 pmol of oligo 5' ccatggagtctgtg (labelled at its 5' end using polynucleotide kinase and ³³P gamma ATP) and buffer and salts according to the suppliers instructions, the total volume being 293 µl. The solution was heated to 65°C and 7 µl of Tth DNA ligase added. The reaction was then cooled to 37°C and left at that 25 temperature for 18hrs. After removal from the reaction solution the strip was washed in T.E. buffer, blotted dry and exposed to a storage phosphor screen from which an image of the radioactivity was taken. The results showed that the target sequence ligated to the correct sequence with a higher yield than to the shorter and longer sequences in adjacent cells of 30 the array.

EXAMPLE 3

Analysis by ligation and polymerisation

- A strip of the array (30mm x 2mm) from Example 2 was added to a solution of 200 pmols of the target oligonucleotide 5' cacagactccatgg(tgaa)₆tgagggaaataag, 200 pmol of oligo 5' ccatggagtctgtg (chemically phosphorylated at the 5' end) with buffer and salts according to the suppliers instructions, the total volume being 243 µl. The solution was heated to 85°C and cooled to 37°C over a period of 30 mins. 7 µl of Tth DNA ligase was added and the reaction mixture heated at 34°C for 17hrs.
- The strip was removed and added to a solution of 8 mM DTT, 3.3 pmol ³²P alpha dTTP, 13 units sequenase version 2.0 and buffer and salts according to the suppliers instructions. The total volume was 250 µl. After heating at 37°C for 3 hrs the strip was removed from the reaction solution, washed in TE buffer, blotted dry and exposed to a storage phosphor screen from which an image of the radioactivity was taken. The results showed counts equal to background over the area of the array where repeats were equal in length or greater than the repeat length of the target, with 20 times the signal in the areas where the repeat length of the array was shorter than the repeat length in the target.

EXAMPLE 4

Analysis by polymerisation

- Two types of polymerase analysis were carried out where reporter nucleotides were chosen, in one case to identify the correct repeat length, and in the other case to identify shorter repeat lengths. This is made possible when the repeat sequence comprises less than all four bases.

- In the former case a base is chosen which is present in the repeat sequence and is different from the first base in the flanking sequence. In the latter case a base is chosen to be complementary to the

first base in the flanking sequence which is absent from the repeat.

A strip of the array from Example 2 (30mm x 2mm) was added to a solution of 500 pmols of the target oligonucleotide 5' cacagactccatgg(tgaa)₆tgagggaaataag in buffer and salts at a concentration 1.09 times the suppliers instructions, the total volume being 275 µl. The solution was heated to 75°C for 5 minutes and cooled to 37°C over a period of 25mins. The solution was removed and added to 3.3 pmols ³²P alpha dCTP, 5 µl 1M DTT. 13 units of Sequenase version 2.0 and water to give a final volume of 295 µl. This solution was added to the array and heated at 37°C for 15hrs 40mins. The polypropylene strip was removed, washed in water and exposed to a storage phosphor screen. The results showed counts of 5 times more for the correct sequence than shorter sequences and twice for the correct sequence compared with longer repeats.

A similar strip of the array (30mm x 2mm) was added to a solution of 500 pmols of the target oligonucleotide 5' cacagactccatgg(tgaa)₆tgagggaaataag in buffer and salts 1.09 times the suppliers instructions, the total volume being 275 µl. The solution was heated to 75°C for 5 mins and cooled to 37°C over a period of 25 mins. The solution was removed and added to 3.3 pmols ³²P alpha dTTP, 5 µl 1M DTT, 13 units of Sequenase version 2.0 and water to give a final volume of 295 µl. This solution was added to the array and heated at 37°C for 15hrs 40mins. The polypropylene strip was removed, washed in water and exposed to a storage phosphor screen. The results showed counts of 4.5 times more for the shorter array sequences than the correct and longer repeat lengths.

EXAMPLE 5

VNTR analysis by ligation

In an experiment similar to the one described in Example 1, an array was created using the human fes/fps locus sequence as a target.

The anchor sequence 5' agagatgtagtctcattcttcgccaggctgg 3' was the actual flanking sequence to the attt repeats of the fes/fps microsatellite (EMBL Accession No X06292 M14209 M14589) as it occurs in human genomic DNA. Using a target oligonucleotide representing the 10 repeat allele and ligating a ³³P labelled 5' flanking sequence (5' g gag aca agg ata gca gtt c 3') and doing a similar experiment to that described above, the resulting radioactivity on the anchor-repeat₁₀ cell was over 10 fold that on adjacent cells in the array.

10

EXAMPLE 6**Demonstration of stepwise ligation of oligonucleotides bound to a solid support.**

A primer oligodeoxynucleotide - 5' PO₄ gta aaa cga cgg cca gt 3', attached to aminated polypropylene through its 3' end, was synthesised and phosphorylated as described. A small square (2mm x 2mm) piece of this material was placed in standard ligation buffer, with template oligonucleotide 5'tcg ttt tac cgt cat gcg tcc tct ctc 3' (250 nM) and a protected ligator oligonucleotide 5' NB PO₄ cgc atg acg 3' (250 nM) and ³³P labelled extender oligonucleotide 5' gag aga gga 3', where NB is a protecting group based on a photocleavable o-nitrobenzyl derivative. The NB protected phosphate of the ligator oligonucleotide had previously been shown to be unable to take part in the ligation reaction. The NB group had also been shown to be removable by uv light to leave a fully functional phosphate group. To this mixture was added thermus thermophilus DNA ligase (Advanced Biotechnologies) 25u and the reaction incubated at room temperature for 6 hours. The mixture was then irradiated with uv light (20 minutes room temperature) and incubated for a further 12 hours. The polypropylene patch was then washed with 30% formamide at 95°C for 5 minutes, and exposed to a storage phosphor screen for 24 hours and scanned in a phosphorimager. The patch showed

a level of radioactivity 50 fold higher than a patch treated in a similar fashion but without addition of the central "ligator" oligonucleotide. In a similar experiment using a phosphorylated ligator oligonucleotide a similar amount of radioactive extender oligonucleotide became covalently
5 attached to a third polypropylene/oligonucleotide primer square.

EXAMPLE 7

Demonstration of point mutation analysis by ligation to allele specific 10 **oligonucleotides attached to solid supports**

Four tethered ASOs 5' (gca or t)ag aga gga 3', differing only at their 5' base, were synthesised as described above, with the 3' end attached to aminated polypropylene. Phosphorylation was carried out as described and four squares of polypropylene carrying each ASO were
15 placed in standard ligation buffer along with complementary target oligonucleotide 5' tcc tct ctc cgt cat gcg tat cgt tca at 3' (250 nM). After addition of ³³P labelled ligator oligonucleotide 5' cgc atg acg 3' (10 nM) and thermus thermophilus DNA ligase (100u), the mixture was incubated at 37°C for 18 hours. The ASO which was fully complementary to the target
20 oligonucleotide was found to have acquired 100-fold greater radioactivity through ligation of the labelled ligator than the non-complementary ASOs.

EXAMPLE 8

Demonstration of DNA ligation specificity

25 In a model experiment to assess the specificity of TTh DNA ligase, ligator and extender deoxyoligonucleotides were ligated together by means of hybridisation to an oligonucleotide template and ligation by DNA ligase.

Template oligonucleotide 5' tcc tct ctc cgt cat gcg tat cgt tca
30 at 3' (250 nM), phosphorylated, ³³P labelled, extender oligonucleotide

5' PO₄gag aga gga 3' (10 nM) and ligator sequence 5' gca gta cg 3' (250 nM) were mixed together in standard ligation buffer with DNA ligase 25 u. This mixture was incubated at 35°C. Samples of this mixture were removed and the reaction stopped by addition of formamide at 15, 30, 60, 5 120, and 240 minutes. The ligated and unligated products were separated by 20% denaturing polyacrylamide gel electrophoresis. The gel was exposed to a phosphor screen for 18 hours and scanned by a phosphorimager. The relative proportions of ligated to unligated products of the reaction were then measured. 50% of the extender sequence had 10 been ligated to the ligator sequence in 30 minutes. By comparison in a similar experiment with ligator 5' gca tga ag 3' after 30 minutes only 1% of the extender sequence had become ligated.

Other polymerases and ligases such as Taq polymerase, Thermosequenase, T4 DNA ligase and *E. Coli* DNA ligase have also been 15 shown to be useful in experiments similar to those described above.

REFERENCES

1. Beaucage, S. L. and Iyer, R. P. (1992). Advances in the synthesis of oligonucleotides by the phosphoramidite approach. 20 Tetrahedron **48**: 2223-2311.
2. Conner, B. J., Reyes, A. A., Morin, C., Itakura, K., Teplitz, R. L., and Wallace, R. B. (1983). Detection of sickle cell β^s globin allele by hybridization with synthetic oligonucleotides. Proc. Natl. Acad. Sci. USA **80**: 278-282.
- 25 3. Cooper, D. N., and Krawczak, M. (1989). The mutational spectrum of single base-pair substitutions causing human genetic disease: patterns and predictions. Hum. Genet. **85**: 55-74.
4. Cotton, RG, (1993) Current methods of mutation detection. Mutation Research **285**: 125-144.

5. Horn, T., and Urdea, M. (1986) Chemical phosphorylation of oligonucleotides. *Tetrahedron Letters* **27**: 4705.
6. Khrapko, K. R., Lysov, Yu. P., Khorlyn, A. A., Shick, V. V., Florentiev, V. L., and Mirzabekov. (1989). An oligonucleotide hybridization approach to DNA sequencing. *FEBS Lett.* **256**: 118-122.
7. Krawczak M. and Schmidtke, J. (1994). DNA fingerprinting. BIOS Scientific Publishers.
8. Maskos, U., and Southern, E.M., (1993) A novel method for the analysis of multiple sequence variants by hybridisation to oligonucleotides. *Nucleic Acids Research*, **21**: 2267-2268.
9. Pillai, V. N. R.(1980). Photoremovable protecting groups in organic chemistry *Synthesis* **39**: 1-26.
10. Southern, E. M. (1988). Analyzing Polynucleotide Sequences. International Patent Application PCT/GB89/00460.
11. Southern, E.M., Maskos, U. and Elder, J.K. (1992). Analysis of Nucleic Acid Sequences by Hybridization to Arrays of Oligonucleotides: Evaluation using Experimental Models. *Genomics* **12**: 1008-1017.
12. Southern, E. M., Case-Green, Elder, J.K. Johnson, M., Mir, K.U., Wang, L., and Williams, J.C. (1994). Arrays of complementary oligonucleotides for analysing the hybridisation behaviour of nucleic acids. *Nucleic Acids Res.* **22**:, 1368-1373.
13. Stahl, S., Hultman, T., Olsson, A., Moks, T.D and, Uhlen, M. (1988) Solid phase DNA sequencing using the biotin-avidin system. *Nucleic Acids Res.* **16**: 3025-38.
14. Veerle, A.M.C.S., Moerkerk, P.T.M.M., Murtagh, J.J., Jr., Thunnissen, F.B.J.M (1994) A rapid reliable method for detection of known point mutations: Point-EXACCT. *Nucleic Acids Research* **22**: 4840-4841.

15. Virnekas, B., Liring, G., Pluckthorn, K., Schneider, C.,
Wellhofer, G. and Moroney, S. E. (1994) Trinucleotide phosphoramidites:
ideal reagents for the synthesis of mixed oligonucleotides for random
mutagenesis. *Nucleic Acids Research* 22: 5600-5607.
- 5 16. Willems, P.J. (1994) Dynamic mutations hit double figures.
Nature Genetics 8: 213-216.

DNA sequences of triplet repeats

Condition	Repeat	Normal	Preexpansion	Expanded	Reference
FRAXA	CGG/CCG	10-50	38-50	200-1000	Verkerk et al. 1991
FRAXE	CGG/CCG			200-1000	Knight et al. 1993
FRAXF	GCC/CGG	6-18	?	300-500	Parrish et al. 1994
FRAX16A	CGG/CCG			1000-2000	Nancarrow et al. 1994
SBMA	CAG	11-31	?	40-62	Tilley et al. 1994
Huntington	CAG	11-34	30-34	42-100	Huntington group 1993
SCA1	CAG	25-36	?	43-81	Orr et al. 1993
DRPLA/HRS	CAG			≤100	Burke et al. 1994
Machado-Joseph	CAG/CTG	~26	?	68-79	Kawaguchi et al. 1994

CLAIMS

1. A method of analysis which comprises: providing a
5 polynucleotide target including a nucleotide at a specified position, and an oligonucleotide probe, tethered to a support, said probe being complementary to the target and terminating at or close to the said specified position; and performing the steps:
 - a) incubating the target with the probe to form a duplex,
 - 10 b) incubating the duplex under ligation conditions with a labelled oligonucleotide complementary to the target,
 - c) and monitoring ligation in b) as an indication of a point mutation at the specified position in the target.
2. A method of analysis which comprises: providing a
15 polynucleotide target having a variable number tandem repeat section and a flanking section, and an oligonucleotide probe having a section complementary to the repeat section and a flanking section of the target; and performing the steps:
 - a) incubating the target with the probe to form a duplex,
 - 20 b) incubating the duplex with a labelled oligonucleotide and/or at least one labelled nucleotide under chain extension conditions,
 - c) and monitoring chain extension as an indication of the length of the variable number repeat section of the target.
3. A method as claimed in claim 2, wherein the probe is
25 tethered to a support.
4. A method as claimed in claim 2 or claim 3, wherein a polymerase and/or ligase is used in step b).

5. A method as claimed in any one of claims 2 to 4, wherein the polynucleotide target has a variable number tandem repeat section and two flanking sections, and wherein in step b) a labelled oligonucleotide is ligated to the probe strand of the duplex.
- 5 6. A method as claimed in any one of claims 2 to 4, wherein the polynucleotide target has a variable number tandem repeat section and two flanking sections, and wherein in step b) the probe strand of the duplex is chain extended by addition of labelled nucleotides.
7. A method as claimed in any one of claims 1 to 6, wherein two
10 or more different probes are tethered to different locations of a support in the form of an array.
8. An array of oligonucleotides, for analysing a polynucleotide target containing a variable sequence, in which each component oligonucleotide i) comprises a sequence complementary to the target
15 including an expected variant of the target, and ii) is tethered to a solid support in a chemical orientation which a) permits duplex formation with the target, and b) permits chain extension only when the sequence of the oligonucleotide matches the variable sequence of the target.
9. A set or array of oligonucleotides, for analysing a
20 polynucleotide target containing a variable number tandem repeat sequence, in which each component oligonucleotide i) comprises a sequence complementary a part of the target immediately adjacent the repeat sequence, ii) comprises a sequence complementary to the repeat sequence of the target and containing a number of repeats expected in the
25 target, and iii) is configured in a way that a) permits duplex formation with the target, and b) permits chain extension only when the number of repeats in the oligonucleotide equals or is less than the number of repeats in the target.

10. A set or array as claimed in claim 9, wherein each component oligonucleotide iii) is configured in a way that b) permits chain extension: by ligation only when the number of repeats in the oligonucleotide equals the number of repeats in the target; or by polymerisation only when the
5 number of repeats in the oligonucleotides is less than the number of repeats in the target.
11. A set or array of oligonucleotides as claimed in claim 9 or claim 10, wherein each component oligonucleotide is tethered to a solid support.
- 10 12. An array of oligonucleotides in which different oligonucleotides occupy different locations and each oligonucleotide has a 3' nucleotide residue through which it is covalently tethered to a support and a 5' nucleotide residue which is phosphorylated.
13. A method of making an array of different oligonucleotides
15 tethered to different locations of a support, which method comprises the steps of: providing a first intermediate oligonucleotide tethered to the support and a second intermediate oligonucleotide in solution, and a third oligonucleotide that is complementary to both the first and second intermediate oligonucleotides, forming a duplex of the third oligonucleotide
20 with the first and second intermediate oligonucleotides, and ligating the first intermediate oligonucleotide with the second intermediate oligonucleotide; and repeating the steps with oligonucleotides tethered to different locations of the support.

Fig.1

Detection of point mutation by single base extension





Fig.2


Detection of point mutation by hybridisation to ASOs
and chain extension



3/7

Fig.3

A. Detection of point mutation by tag ligation to ASOs



B. Detection of point mutation by ligation to library of differentially tagged ASOs






Fig.4a

Analysis of VNTRs by ligation of tag to allelic variants


A. Target repeats equal probe



B. Target repeats more than probe



C. Target repeats less than probe




5/7


Fig.4b

Analysis of VNTRs by ligation of tag to allelic variants

A. Target repeats equal probe



B. Target repeats more than probe



C. Target repeats less than probe





Fig.5

Measurement of VNTRs by labelled chain extension




7/7


Fig.6

Analysis of VNTRs by ligation of tag followed by chain extension


A. Target repeats equal probe



B. Target repeats more than probe



C. Target repeats less than probe



SUBSTITUTE SHEET (RULE 26)

INTERNATIONAL SEARCH REPORT

Internat Application No
PCT/GB 96/00848

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C12Q1/68

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 C12Q

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO,A,95 09248 (ARCH DEVELOPMENT CORP) 6 April 1995 see page 53, line 5 - page 54, line 2 ---	1-13
Y	WO,A,94 11530 (TRUSTEES OF BOSTON UNIVERSITY) 26 May 1994 see the whole document ---	1-13
Y	WO,A,95 04160 (ISIS INNOVATION LTD17) 9 February 1995 see page 18, line 27 - line 30 ---	1-13
Y	WO,A,93 25563 (CITY OF HOPE) 23 December 1993 see the whole document ---	1-12
-/--		

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

13 August 1996

Date of mailing of the international search report

03.09.96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+ 31-70) 340-3016

Authorized officer

Osborne, H

INTERNATIONAL SEARCH REPORT

Internat. Application No.

PCT/GB 96/00848

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	EP,A,0 246 864 (IMPERIAL CHEMICAL INDUSTRIES PLC) 25 November 1987 see claims 1-14 ---	1-12
Y	EP,A,0 552 545 (PIONEER HI-BRED INTERNATIONAL, INC.) 28 July 1993 see the whole document ---	1-12
A	EP,A,0 185 494 (APPLIED BIOSYSTEMS INC) 25 June 1986 see the whole document ---	1-12
P,X	WO,A,95 27078 (CIBA CORNING DIAGNOSTICS CORP) 12 October 1995 see the whole document ---	1-12
A	TRENDS IN BIOTECHNOLOGY, vol. 11, June 1993, pages 238-46, XP000440486 PROSSER J.: "detecting single base mutations" see page 242, column 2 - page 245 -----	1-13

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GG 96/00848

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9509248	06-04-95	AU-B- 8072794 CA-A- 2172722 EP-A- 0723598 NO-A- 961165	18-04-95 06-04-95 31-07-96 23-05-96
WO-A-9411530	26-05-94	EP-A- 0668932 US-A- 5503980	30-08-95 02-04-96
WO-A-9504160	09-02-95	AU-B- 7269194 CA-A- 2168010 EP-A- 0711362 FI-A- 960403 NO-A- 960370	28-02-95 09-02-95 15-05-96 29-01-96 28-03-96
WO-A-9325563	23-12-93	CA-A- 2115342 AU-B- 2251192 EP-A- 0607151 JP-T- 6509946	23-12-93 04-01-94 27-07-94 10-11-94
EP-A-246864	25-11-87	AT-T- 108490 DE-D- 3750198 DE-T- 3750198 ES-T- 2058112 HK-A- 125195 JP-A- 63022197	15-07-94 18-08-94 26-01-95 01-11-94 11-08-95 29-01-88
EP-A-552545	28-07-93	AU-B- 3107493 CA-A- 2085559 HU-A- 69947 JP-A- 7246099 NZ-A- 245242	22-07-93 18-07-93 28-09-95 26-09-95 22-12-94
EP-A-185494	25-06-86	US-A- 4883750 DE-A- 3586090 JP-B- 6044880 JP-A- 61191964 US-A- 5521065 US-A- 5242794	28-11-89 25-06-92 15-06-94 26-08-86 28-05-96 07-09-93
WO-A-9527078	12-10-95	AU-B- 1958595	23-10-95